

Published with Open Access at **Journal BiNET**

Vol. 02, Issue 02: 88-100

Journal of Fisheries, Livestock and Veterinary ScienceJournal Home: <https://www.journalbinet.com/jflvs-journal.html>

Stress biomarker: A review on glucocorticoids concentration pattern & its impact on captive animals

Purba Islam¹, Pritam Saha¹, Popy Khatun¹, Md. Imran Hossain¹, Arup Islam² and Sabhya Sachi¹

¹Department of Pharmacology, Faculty of Veterinary Science, Bangladesh Agricultural University, Bangladesh

²Department of Microbiology, Mymensingh Medical College, Bangladesh

✉ For any information: purba.islam@bau.edu.bd (Islam, P)

Article received: 13.05.22; Revised: 28.07.22; First published online: 10 August, 2022.

ABSTRACT

As a stress biomarker, glucocorticoids (GCs) concentration in wild animals shows how well a species copes with confinement. In captivity, the animal may suffer from many health problems such as weight loss, irreversible glucocorticoid changes, immune system abnormalities, and reproductive suppression. The review mainly focuses on the effect of transferring wild-caught animals to captivity on stress-related physiological systems such as weight changes, glucocorticoid modulation, adrenomedullary control, and effects on immune and reproductive systems. The GCs concentration pattern on species-wise variation is recorded, and the detection technique and alleviation process is also discussed.

Key Words: Captivity, Stress, Glucocorticoids, Impacts, Detection technique and Alleviation.

Cite Article: Islam, P., Saha, P., Khatun, P., Hossain, M. I., Islam, A. and Sachi, S. (2022). Stress biomarker: A review on glucocorticoids concentration pattern & its impact on captive animals. *Journal of Fisheries, Livestock and Veterinary Science*, 02(02), 88-100.

Crossref: <https://doi.org/10.18801/jflvs.020222.10>



Article distributed under terms of a Creative Common Attribution 4.0 International License.

I. Introduction

Stress is a condition that endangers or appears to endanger the homeostasis of an individual, and in response to stress, an individual adapts a series of mechanisms aimed at protecting and restoring homeostasis (Möstl and Palme, 2002; Sapolsky, 2000). Stress responses play a crucial role in allowing animals to cope with change and challenges to face both environmental certainty and uncertainty (Sheriff et al., 2011). The goal of keeping wild animals in captivity is for conservation, research, agriculture, the exotic pet trade, and entertainment (Morgan and Tromborg, 2007). Animals kept in captivity have a comfortable home and plenty of food. Although the basic needs of animals are met in captivity, the conditions of confinement can result in physiological stress, especially in newly captured animals (Fischer and Romero, 2019). Human presence, an unfamiliar environment, artificial light conditions, confinement, dietary changes, and handling of animals are all unpredictable and uncontrollable inputs that may activate the stress response (Morgan and Tromborg, 2007). The behaviour and adrenal activity of animals are also influenced by visitors (Pifarre et al., 2012).

The stress response is mediated by several hormones, including ACTH, glucocorticoids, catecholamines, prolactin, and others (Moberg, 2000). In stress conditions, glucocorticoid hormones, mainly cortisol, are secreted by the adrenal glands and regulate energy and maintain homeostasis (Sapolsky et al., 2000). Increased plasma levels of glucocorticoid hormones like corticosterone are critical physiological reactions to stressful events in vertebrates (Moore and Jessop, 2003; Romero, 2004). Most taxa, including birds (Cockrem and Silverin, 2002), reptiles (Cash et al., 1997), mammals (Lopez-Olvera et al., 2007), and amphibians (Mosconi et al., 2006), exhibit this response. Cortisol levels that are too high can impact the behaviour of animals (O'Connor et al., 2000). In wild animals, short-term captivity is known to cause a hormonal stress response (Mosconi et al., 2006). In long term captivity, the increased level of cortisol may influence behaviour (more aggression, moving, and grooming activities) and physiological functions such as depression, hypertension, loss of body weight, reproductive failure, immunosuppression, and a shorter life span (Birke, 2002; Breazile, 1987; Palme, 1997).


To keep the animals healthy, stress symptoms should be decreased or eliminated. In mammals, the relationship between stress hormones and leukocytes has been extensively established (Dhabhar et al., 1994). Ornithologists are increasingly using the neutrophil/heterophil ratio to demonstrate the consequences of various stressful conditions, such as transit stress (Groombridge et al., 2004) and reproductive output (Moreno et al., 2002). The success of modern zoos and safaris is dependent on achieving and maintaining high animal care standards. Therefore, stress levels should be eliminated, and better management should be provided to ensure animal welfare. The review documented the effects of stress on captive animals, especially the patterns of change in baseline and integrated GCs. Finally, some mitigation process is discussed to improve the condition of captive animals.

II. Materials and Methodology

Published literature on glucocorticoids as a stress indicator was collected from PubMed, Scientific Gate, Google Scholar, and E-Journals of ISI (Institute for Scientific Information). The following keywords were used to search the literature: "glucocorticoids," "captivity," "stress," "physiology," or "endocrinology," and related words. The searched items or publications were thoroughly checked, downloaded in detail, and reviewed. Only the original research data containing publications written in English were included for our review. The abstracts of the research articles that had data regarding our interests were selected for review of the full content. The research work was thoroughly revised and sorted out to meet the field of interest. The full articles were managed in PDF format using Mendeley, a reference management software. We, therefore, devised the following criteria to determine whether papers should be included: (a) GCs elevated in captivity (b) GCs increase at capture then decrease (c) GCs lower in captivity (d) High initial GCs decrease after the captivity (e) No effect of GCs (f) Impact of stress on captive animal

Publications

Primarily, we found 210 articles that could meet the area of interest. A total of 140 articles contained original research data, and the English language was used to write up to 125. Out of 125, we considered 56 reports for our study due to a lack of detailed information in the remaining papers (both quantitative and qualitative) on glucocorticoid concentration in captive wild animals.



Species-wise analysis

Among the surveyed species, the highest number of works, 15 (40.54%), were accomplished on mammals, followed by birds, 9 (24.32%); reptiles, 7 (18.91%) and amphibians, 6 (16.21%) (Table 01). It is observed that more research studies on glucocorticoid concentration were performed on mammalian species than on other animals (Figure 01).




Figure 01. Research on GCs (Species-wise analysis)

III. Glucocorticoids level in wild animals

Under stress conditions, GCs are secreted from the adrenal glands and cause many health problems to the animal. So GCs can be a valuable source of knowing the stress level in animals. The review documented the data about the species, the duration of the study, the sample used for GC determination, ranges of GC concentration, as well as the GC concentration in different conditions to know the worldwide scenario of the stress level in animals.

Table 01. Patterns of change in baseline and integrated GCs when wild animals are brought into captivity

Mammals						
Species	Study duration	Sample used for GC determination	Detection technique	Detection ranges	GC concentration pattern in captivity	Normal level of cortisol
Canada lynx (<i>Lynx canadensis</i>) (Fanson et al., 2012)	Long term (unknown)	Faecal Glucocorticoids Materials (FGMs)	EIA	350 ng/ml		25 ng/ml
Spider monkey (<i>Ateles geoffroyi yucatanensis</i>) (Rangel-Negrín et al., 2009)	Long term (unknown)	FGMs	RIA	1925.4 ng/g		1224.2 ng/g
Grevy's zebra (<i>Equus grevyi</i>) (Franceschini et al., 2008)	6 weeks	FGMs	RIA	25.1 ng/gm	Elevated	14.6 ng/gm
Black rhinos White rhinos (<i>Ceratotherium simum</i>) (Linklater et al., 2010; Yang et al., 2019)	75 days	FGMs	RIA	Black Rhino (Male 23.9 ng/gm; Female 24.5 ng/gm) White Rhino (Male 12.3 ng/gm; Female 16.3 ng/gm)		9.34 ng/gm
Degu (<i>Octodon degus</i>) (Quispe et al., 2014)	>1 year	Plasma	RIA	150 ng/ml		100ng/ml
Harbor seal (<i>Phoca vitulina</i>) (Trumble et al., 2013)	>4 weeks	Plasma	RIA	125 ng/ml	No effect	148 ng/ml

Mammals						
Species	Study duration	Sample used for GC determination	Detection technique	Detection ranges	GC concentration pattern in captivity	Normal level of cortisol
Tuco-tuco (<i>Ctenomys talarum</i>) (Vera et al., 2011)	30 days	Plasma	RIA	18.97ng/ml		14.40 ng/ml
Chacma baboon (<i>Papio ursinus</i>) (Steyn, 1975)	Peak: 4 weeks approach long term captives by 7 weeks	Plasma	RIA	141 mg/100 ml	Increase at capture, then decrease to approach wild baseline	99.90 mg/100 ml
African green monkey (<i>Cercopithecus aethiops</i>) (Suleman et al., 2004)	Peak: 1 day approach free-living by 2 days	Plasma	RIA	530 nmol/L		391 nmol/L
Richardson's ground squirrel (<i>Urocyon richardsoni</i>) (Hare et al., 2014)	Peak: 3-5 days approach free-living by 6 days	FGMs	RIA	161.04 ng/ml		104.99 ng/ml
Harbor porpoise (<i>Phocoena phocoena</i>) (Siebert et al., 2011)	Long term	Plasma	RIA	122 µg/L		338.6 µg/L
Gilbert's potoroo (<i>Potorous gilbertii</i>) (Stead-Richardson et al., 2010)	Long term	FGMs	RIA	6 ng/ml	Lower	10.5 ng/ml
Harbor seal (<i>Phoca vitulina</i>) (Gardiner and Hall, 1997)	Long term	Plasma	ELISA	444.2 nmol/l		829.5 nmol/l
Bushtail possums (<i>Trichosurus volpecula</i>) (female) (Baker et al., 1998)	Decreased from week 1 to 20	Plasma	RIA	34 ng/ml	Elevated at wild baseline decrease over capture period	13 ng/ml
Meadow vole (<i>Microtus pennsylvanicus</i>) (Olsen and Seabloom, 1973)	Decreased from day 1 to day 70	plasma	Fluorometric Determination	Male 1397 ng/ml Female 1562 ng/ml		Male 300 ng/ml Female 590 ng/ml

Birds						
Species	Study Duration	Sample used for GC determination	Detection technique	Detection ranges	GC concentration pattern in captivity	Normal level of cortisol
White crowned sparrow (<i>Zonotrichia leucophrys</i>) (Marra et al., 1995)	35 days	Plasma	RIA	16.45 ng/ml		13.25 ng/ml
Blackbirds (<i>Turdus merula</i>) (Adams et al., 2011)	22 days	Plasma	RIA	6.8 ng/ml	Elevated	2.2 ng/ml
House sparrow (<i>Passer domesticus</i>) (Fischer and Romero, 2016)	7 days	Plasma	RIA	4.8 ng/ml		1.2 ng/ml
Southern pied babbler (<i>Turboides bicolor</i>) (Jepsen et al., 2012)	5 days	FGMs	EIA	73.49 ng/ml		21.46 ng/ml
House sparrow (<i>Passer domesticus</i>) (Martin et al., 2011)	Up to 4 weeks	Plasma	ELISA	20 ng/ml	No effect	1.2 ng/ml

Birds						
Species	Study Duration	Sample used for GC determination	Detection technique	Detection ranges	GC concentration pattern in captivity	Normal level of cortisol
House sparrow (<i>Passer domesticus</i>) (Kuhlman and Martin, 2010)	Peak: Days 1-2 approach free-living by 1 month	Plasma	EIA	20 ng/ml	Increase at capture, then decrease to approach wild baseline	1.2 ng/ml
White crowned sparrow (<i>Zonotrichia leucophrys</i>) (Wingfield et al., 1982)	Peak: Days 1-2 approach free-living by day 14	Plasma	RLA	25.39 ng/ml		15.77 ng/ml
European starling (<i>Stumus vulgaris</i>) (Cyr and Romero, 2008)	Unknown	FGMs	RIA	7 ng/ml	Lower	55 ng/ml
Red knot (<i>Calidris canutus</i>) (Piersma and Ramenofsky, 1998; Piersma et al., 2000)	Decreased from first sample to 2 years	Plasma	RIA	30 ng/ml	Elevated at wild baseline decrease over capture period	40ng/ml
Reptiles						
Species	Study Duration	Sample used for GC determination	Detection technique	Detection ranges	GC concentration pattern in captivity	Normal level of cortisol
Tuatara (<i>Sphenodon punctatus</i>) (female) (Tyrrell and Cree, 1994)	Unknown	Plasma	RIA	Captive juvenile female 4.21 ng/ml Wild juvenile female 2.44 ng/ml	Elevated	1.28-4.65 ng/ml
Tree lizard (<i>Urosaurus ornatus</i>) (Moore et al., 1991)	Up to 3 weeks	Plasma	RIA	20 ng/ml		10ng/ml
Water snake (Nerodia sipedon) (Sykes and Klukowski, 2009)	5-8 days	Plasma	ELISA	310 ng/ml	Elevated	275 ng/ml
Brown tree snake (<i>Boiga irregularis</i>) (Mathies et al., 2001)	3 days	Plasma	RIA	Male 49.30 ng/ml Female 56.87 ng/ml		Male 4.41 ng/ml Female 8.71 ng/ml
Tuatara (<i>Sphenodon punctatus</i>) (male) (Tyrrell and Cree, 1994)	Unknown	Plasma	RIA	3.61 ng/ml	No effect	2.48 ng/ml
Kutum (<i>Rutilus frisii kutum</i>) (Nikoo et al., 2010)	3 days	Plasma	RIA	308 ng/ml		378 ng/ml
Skink (<i>Egernia whitii</i>) (Jones and Bell, 2004)	Peak: 1 day-1 week approach free-living by 4 weeks	Plasma	RIA	24.8 ng/ml	Increase at capture, then decrease to approach wild baseline	5.3 ng/ml

Amphibians						
Species	Study Duration	Sample used for GC determination	Detection technique	Detection ranges	GC concentration pattern in captivity	Normal level of cortisol
Cururu toad (<i>Rhinella icterica</i>) (De Assis et al., 2015)	3 months	Plasma	EIA	65 ng/ml	Elevated	9 ng/ml
Toad (<i>Rhinella schneideri</i>) (Titon et al., 2017; Narayan et al., 2012)	60 days	Plasma	EIA	4.24 ng/ml		31.3 pg/ μ g
Cane toad (<i>Rhinella marina</i>) (Narayan and Hero, 2011)	Peak: Day 5 approach free-living by Day 12	Urine	RIA	600 pg/ μ g	Increase at capture, then decrease to approach wild baseline	200 pg/ μ g
Cane toad (<i>Rhinella marina</i>) (Narayan et al., 2012)	Peak: Day 4 approach free-living by Day 14	Urine	EIA	66.3 pg/ μ g		31.3 pg/ μ g
Fijian ground frog (<i>Platymanthis vitiana</i>) (Narayan and Hero, 2011)	Peak: Day 5 approach free-living by Day 25	Urine	EIA	113.93 pg/ μ g		34.74 pg/ μ g
Tod (<i>Rhinella icterica</i>) (Titon et al., 2018)	Decreased from days 30 to 60	Plasma	RIA	3ng/ml	Lower	18 ng/ml

IV. Glucocorticoids concentration pattern-wise analysis

Most of the research findings have shown that elevated glucocorticoids concentrations at 14 (37.83%) in captivity, followed by increased glucocorticoids at capture, then decreased to approach wild baseline, 8 (21.62%), the same percentage of lower and no effect on glucocorticoid concentration in captivity, 6 (16.21%), and then elevated glucocorticoids concentrations at wild baseline decreased over the capture period, 3 (8.1%) (Figure 02).




Figure 02. Patterns of change in baseline and integrated GCs when wild animals are brought into captivity

V. Detection technique

GCs concentrations are one of the most commonly utilized measures to assess captivity stress. Acute stress causes a brief increase in GCs. Long-term stressor exposure, on the other hand, typically causes alterations in GC regulation that vary depending on the species and conditions (Dickens and Romero, 2013). Thus monitoring glucocorticoid levels can be a valuable tool to provide information on the stress level experienced by the animals. Glucocorticoid levels can be found in blood, faeces, hair, etc. (Cockrem, 2005; Goymann et al., 1999). The most frequent method for measuring GCs is quantifying them in circulating plasma (Sheriff et al., 2011). Nowadays, the assessment of hormone matrix using

faeces is a widely accepted approach. The benefit of faeces collection is, easy to collect, animals are usually not disturbed during sample collection, and sampling is feedback-free due to the absence of capture and handling (Ganswindt et al., 2012).

Enzyme-Linked ImmunoSorbent Assay (ELISA), Chemiluminescent Immuno Assays (CLIA), Radio Immuno Assays, mass spectrometry (MS), and other immunoassays are the most commonly used for detection and quantification of cortisol.

In immunoassays, antibodies are used to produce a signal and subsequently determine the concentration of the target compound (i.e., the analyte); in an ELISA, one of the employed antibodies is modified (enzyme-conjugated) to be able to produce an optical signal after administration of the substrate; in a competitive CLIA, the target compound (e.g., cortisol or cortisone) is labeled by chemiluminescent tracer substance (Hogenelst et al., 2019). When a specific antibody is added to the labelled antigen, they will bind to each other. Another immunoassay method (LFIA) is the Lateral Flow Immuno Assay, typically employed for qualitative evaluations (Posthuma-Trumpie et al., 2009). In particular, a liquid is placed on a strip and moved over a test strip through several compartments as it passes over a liquid (such as urine or saliva). As a result, a coloured strip will appear if the analyte is present.

Mass-spectrometry (MS) is a technique that separates compounds based on their polarity and detects them based on their mass and charge. Because of its great specificity and sensitivity, it is regarded as the gold standard for cortisol measurement (Miller et al., 2013; Perogamvros et al., 2010; Baid et al., 2007).

The Radio Immune Assay (RIA) and High-Performance Liquid Chromatography are the common detection techniques to measure faeces' GCs levels (Ganswindt et al., 2012).

VI. Impact of elevated glucocorticoid concentration on animals

GC concentrations were one of the most common approaches to quantifying captivity stress. GC hormones (mostly cortisol in fish and most mammals; primary corticosterone in reptiles, birds, amphibians, and rodents) are produced by the adrenal cortex and have a variety of actions throughout the body (Fischer and Romero, 2019). Acute stresses cause a spike in GCs that is quickly controlled by negative feedback. Changes in GC regulation are common after long-term stressor exposure. But depending on the species and the environment, different parts of the GCs response (baseline concentrations, stress-induced concentrations, or negative feedback) and the direction of change are impacted (Dickens and Romero, 2013). The GCs of different animals are not affected by captivity. In 16.21% (6 of 37) of the studies, there was no recorded difference in GCs during or after captivity compared to free-living levels. However, in most studies, captivity caused a change in baseline or integrated GCs. In 37.83% of studies (14 of 37), wild animals had increased GCs at the end of the capture period compared to concentrations in free-living animals (3 days to several years). Traditionally, elevated GCs have been considered a sign that animals are persistently stressed. This stress state can have serious consequences for animals, some of which are discussed in this article.

Weight loss

When wild animals are brought into captivity, they frequently lose weight for a time. According to a study of wild animals taken into captivity, 64 % of the animals examined exhibited a decrease in bulk as a result of being imprisoned at least during the first catch phase (Fischer and Romero, 2019). In captivity, chronic stress is most likely to blame for weight loss. Weight loss has been observed in mammals (Flügge, 1996), birds (Rich and Romero, 2005) and fish (Rich and Romero, 2005). Weight loss is the most prevalent symptom of persistent stress (Dickens and Romero, 2013). Weight loss was not the only trend observed in captivity. Fischer and Romero, 2019 found that 17% of the studied animals acquired weight beyond their initial condition during their investigation (Fischer and Romero, 2019). Because of the greater food available in confinement, certain animals may maintain their weight. On the other hand, other animals may grow obese due to unrestricted availability to food and exercise limits and suffer the negative consequences of having a big body mass or body fat content (West and York, 1998). In a study with domesticated budgerigars, birds were fed ad libitum and housed in cages with limited movement, and high body mass at the end of 28 days was linked to greater DNA damage (Larcombe et al., 2015).

Immune consequences

Stress has some effects on the immune system. Increased GCs have a temporary or long-term impact on leukocyte populations, accounting for these changes. Lymphocytes moving from the bloodstream to the skin, spleen, and lymph nodes, where they will be ready in the event of a wound, can occur as a result of GCs (Dhabhar and McEwen, 1997). Neutrophils (most vertebrates) and heterophils (birds and some reptiles) can both multiply and mobilize in response to GCs (Dale et al., 1975; Gross and Siegel, 1983). As a result of these effects on leukocyte populations, the neutrophil or heterophil to lymphocyte ratio (N or H: L ratio) alters (Dhabhar and McEwen, 1997). A change in the N or H:L ratio does not always mean that the immune system of an animal is underactive or overactive. Instead, it serves as a supplementary indicator, similar to GC secretion. Davis et al. (2008) found that a long-term increase in the N or H:L ratio, equal to a long-term increase in circulating GCs, can suggest that an animal is stressed. Overall, it appears that confinement has no distinct immune regulatory pattern. Although it has been shown that captivity reduces immune activity in some animals (e.g., red knots and toads), it has also increased immune function in others. Gene expression for pro-inflammatory cytokines was higher in imprisoned house sparrows (2 and 4-week captives) than in recently acquired animals in house sparrows, indicating that captive birds are more prone to inflammation (Martin et al., 2011). GCs release is most likely linked to changes in the immune system's response to prolonged stress. The impact of GCs on the immune system, on the other hand, can be complicated. GCs often induce an immunological response in the short term, but they may be immunosuppressive in the long run (Dhabhar and McEwen, 1997; Martin, 2009). It is impossible to predict whether captivity conditions will result in appropriate or inappropriate immune activity because the interaction between GCs and immunity is complex and context-specific.

Reproductive function

The adverse effects of captivity on reproductive biology have been well documented. High GC levels are likely linked to a reduced ability to reproduce. GCs may help reduce the negative effects of reproductive hormones (Sapolsky et al., 2000). Long-term GC exposure can influence gonad growth, egg maturation, sperm production, and behaviour by lowering testosterone and estradiol levels. A drop in sex hormones was followed by increased GCs in green tree frogs. (Zerani et al., 1991). Despite GC levels being below free-living values for most of the confined period, males had suppressed faecal testosterone, and females had suppressed faecal progesterins in black rhinos (Linklater et al., 2010).

VII. Alleviation of stress in captivity

One of the simplest methods to minimize chronic stress symptoms is to change the physical conditions of captivity. In European starlings (long term captivity) (Dickens and Bentley, 2014), transferring from outdoor to indoor cages resulted in lower reproductive hormones and behaviours, as well as weight loss and immunological function (Dickens and Bentley, 2014). The size and density of the cage play a role in the development of chronic stress. During the initial captivity period high-density housing resulted in elevated GCs compared to low-density housing (Bolasina, 2011). Adding a female to a cage of male brown-headed cowbirds (who had previously been housed alone) resulted in lower plasma GCs and higher testicular regeneration. For visual species, lighting conditions could be critical. Under low flicker rate fluorescent lights, European starlings display more behavioural symptoms of chronic stress than under high flicker rate fluorescent lights (Evans et al., 2012), but low flicker rate fluorescent lights do not elicit a GC response (Greenwood et al., 2004). Particularly for poikilotherms, temperature conditions should be carefully considered. During the first transfer to captivity, warm conditions resulted in significant mortality in sardines (Marçalo et al., 2008) and increased GCs in cane toads (Narayan et al., 2012).

VIII. Conclusion

Captivity, a new site, travel, social tension and hostility, human disturbances, and predator exposure have all been used to identify adrenocortical activity using GCs. Captivity has been proven to raise baseline GCs and impair adrenal sensitivity in the literature. Even if the animal's basic needs are supplied, high GCs in many species can persist for months or years after capture, resulting in chronic stress. As a result, stress may impact the physical, psychological, and reproductive behaviour of zoo animals. Individual animal factors (such as species traits, genetics, temperament, and previous experience) play a role in how well an animal copes with environmental conditions (e.g., social

grouping, enclosure design, and sensory environment). The most noteworthy finding of this study is that GCs are generally higher in captivity and have a deleterious influence on captive animals.

IX. References

- [1]. Adams, N. J., Farnworth, M. J., Rickett, J., Parker, K. A. and Cockrem, J. F. (2011). Behavioural and corticosterone responses to capture and confinement of wild blackbirds (*Turdus merula*). Applied Animal Behaviour Science, 134(3-4), 246-255. <https://doi.org/10.1016/j.applanim.2011.07.001>
- [2]. Baid, S. K., Sinaii, N., Wade, M., Rubino, D. and Nieman, L. K. (2007). Radioimmunoassay and tandem mass spectrometry measurement of bedtime salivary cortisol levels: a comparison of assays to establish hypercortisolism. The Journal of Clinical Endocrinology and Metabolism, 92(8), 3102-3107. <https://doi.org/10.1210/jc.2006-2861>
- [3]. Baker, M. L., Gemmell, E. and Gemmell, R. T. (1998). Physiological changes in brushtail possums, *Trichosurus vulpecula*, transferred from the wild to captivity. Journal of Experimental Zoology, 280(3), 203-212. [https://doi.org/10.1002/\(SICI\)1097-010X\(19980215\)280:3<203::AID-JEZ1>3.0.CO;2-R](https://doi.org/10.1002/(SICI)1097-010X(19980215)280:3<203::AID-JEZ1>3.0.CO;2-R)
- [4]. Birke, L. (2002). Effects of browse, human visitors and noise on the behaviour of captive orang utans. Animal Welfare Potters Bar, 11(2), 189-202.
- [5]. Bolasina, S. N. (2011). Stress response of juvenile flounder (*Paralichthys orbignyanus*, Valenciennes 1839) to acute and chronic stressors. Aquaculture, 313(1-4), 140-143. <https://doi.org/10.1016/j.aquaculture.2011.01.011>
- [6]. Breazile, J. E. (1987). Physiological basis and consequences of distress in animals. Journal of The American Veterinary Medical Association, 191, 1212-1215.
- [7]. Cash, W. B., Holberton, R. L. and Knight, S. S. (1997). Corticosterone secretion in response to capture and handling in free-living red-eared slider turtles. General and comparative endocrinology, 108(3), 427-433. <https://doi.org/10.1006/gcen.1997.6999>
- [8]. Cockrem, J. F. (2005). Conservation and behavioral neuroendocrinology. Hormones and Behavior, 48(4), 492-501. <https://doi.org/10.1016/j.yhbeh.2005.03.008>
- [9]. Cockrem, J. F., and Silverin, B. (2002). Sight of a predator can stimulate a corticosterone response in the great tit (*Parus major*). General and comparative endocrinology, 125(2), 248-255. <https://doi.org/10.1006/gcen.2001.7749> <https://doi.org/10.1006/gcen.2001.7750>
- [10]. Cyr, N. E. and Romero, L. M. (2008). Fecal glucocorticoid metabolites of experimentally stressed captive and free-living starlings: implications for conservation research. General and Comparative Endocrinology, 158(1), 20-28. <https://doi.org/10.1016/j.ygcen.2008.05.001>
- [11]. Dale, D. C., Fauci, A. S. and Wolff, S. M. (1975). Comparison of agents producing a neutrophilic leukocytosis in man. Hydrocortisone, prednisone, endotoxin, and etiocholanolone. The Journal of clinical investigation, 56(4), 808-813. <https://doi.org/10.1172/JCI108159>
- [12]. Davis, A. K., Maney, D. L., and Maerz, J. C. (2008). The use of leukocyte profiles to measure stress in vertebrates: a review for ecologists. Functional Ecology, 22(5), 760-772. <https://doi.org/10.1111/j.1365-2435.2008.01467.x>
- [13]. De Assis, V. R., Titon, S. C. M., Barsotti, A. M. G., Titon Jr, B., and Gomes, F. R. (2015). Effects of acute restraint stress, prolonged captivity stress and transdermal corticosterone application on immunocompetence and plasma levels of corticosterone on the cururu toad (*Rhinella icterica*). The Public Library of Science, 10(4), e0121005. <https://doi.org/10.1371/journal.pone.0121005>
- [14]. Dhabhar, F. S. and McEwen, B. S. (1997). Acute stress enhances while chronic stress suppresses cell-mediated immunity in vivo: A potential role for leukocyte trafficking. Brain, behavior, and immunity, 11(4), 286-306. <https://doi.org/10.1006/brbi.1997.0508>
- [15]. Dhabhar, F. S., Miller, A. H., Stein, M., McEwen, B. S. and Spencer, R. L. (1994). Diurnal and acute stress-induced changes in distribution of peripheral blood leukocyte subpopulations. Brain, behavior, and immunity, 8(1), 66-79. <https://doi.org/10.1006/brbi.1994.1006>
- [16]. Dickens, M. J. and Bentley, G. E. (2014). Stress, captivity, and reproduction in a wild bird species. Hormones and Behavior, 66(4), 685-693. <https://doi.org/10.1016/j.yhbeh.2014.09.011>
- [17]. Dickens, M. J. and Romero, L. M. (2013). A consensus endocrine profile for chronically stressed wild animals does not exist. General and comparative endocrinology, 191, 177-189. <https://doi.org/10.1016/j.ygcen.2013.06.014>

- [18]. Evans, J. E., Smith, E. L., Bennett, A. T., Cuthill, I. C., and Buchanan, K. L. (2012). Short-term physiological and behavioural effects of high-versus low-frequency fluorescent light on captive birds. *Animal Behaviour*, 83(1), 25-33. <https://doi.org/10.1016/j.anbehav.2011.10.002>
- [19]. Fanson, K. V., Wielebnowski, N. C., Shenk, T. M. and Lucas, J. R. (2012). Comparative patterns of adrenal activity in captive and wild Canada lynx (*Lynx canadensis*). *Journal of Comparative Physiology B*, 182(1), 157-165. <https://doi.org/10.1007/s00360-011-0597-8>
- [20]. Fischer, C. P. and Romero, L. M. (2016). The use of α - or β -blockers to ameliorate the chronic stress of captivity in the house sparrow (*Passer domesticus*). *Conservation Physiology*, 4(1). <https://doi.org/10.1093/conphys/cow049>
- [21]. Fischer, C. P. and Romero, L. M. (2019). Chronic captivity stress in wild animals is highly species-specific. *Conservation physiology*, 7(1), coz093. <https://doi.org/10.1093/conphys/coz093>
- [22]. Flügge, G. (1996). Alterations in the central nervous α 2-adrenoceptor system under chronic psychosocial stress. *Neuroscience*, 75(1), 187-196. [https://doi.org/10.1016/0306-4522\(96\)00292-8](https://doi.org/10.1016/0306-4522(96)00292-8)
- [23]. Franceschini, M. D., Rubenstein, D. I., Low, B. and Romero, L. M. (2008). Fecal glucocorticoid metabolite analysis as an indicator of stress during translocation and acclimation in an endangered large mammal, the Grevy's zebra. *Animal Conservation*, 11(4), 263-269. <https://doi.org/10.1111/j.1469-1795.2008.00175.x>
- [24]. Ganswindt, A., Brown, J. L., Freeman, E. W., Kouba, A. J., Penfold, L. M., Santymire, R. M. and Milnes, M. R. (2012). International Society for Wildlife Endocrinology: the future of endocrine measures for reproductive science, animal welfare and conservation biology. <https://doi.org/10.1098/rsbl.2011.1181>
- [25]. Gardiner, K. J. and Hall, A. J. (1997). Diel and annual variation in plasma cortisol concentrations among wild and captive harbor seals (*Phoca vitulina*). *Canadian Journal of Zoology*, 75(11), 1773-1780. <https://doi.org/10.1139/z97-806>
- [26]. Goymann, W., Möstl, E., Van't Hof, T., East, M. L. and Hofer, H. (1999). Noninvasive fecal monitoring of glucocorticoids in spotted hyenas, *Crocuta crocuta*. *General and Comparative Endocrinology*, 114(3), 340-348. <https://doi.org/10.1006/gcen.1999.7268>
- [27]. Greenwood, V. J., Smith, E. L., Goldsmith, A. R., Cuthill, I. C., Crisp, L. H., Walter-Swan, M. B. and Bennett, A. T. (2004). Does the flicker frequency of fluorescent lighting affect the welfare of captive European starlings? *Applied Animal Behaviour Science*, 86(1-2), 145-159. <https://doi.org/10.1016/j.applanim.2003.11.008>
- [28]. Groombridge, J. J., Massey, J. G., Bruch, J. C., Malcolm, T. R., Brosius, C. N., Okada, M. M. and Sparklin, B. (2004). Evaluating stress in a Hawaiian honeycreeper, *Paroreomyza montana*, following translocation. *Journal of Field Ornithology*, 75(2), 183-187. <https://doi.org/10.1648/0273-8570-75.2.183>
- [29]. Gross, W. B. and Siegel, H. S. (1983). Evaluation of the heterophil/lymphocyte ratio as a measure of stress in chickens. *Avian diseases*, 972-979. <https://doi.org/10.2307/1590198>
- [30]. Hare, J. F., Ryan, C. P., Enright, C., Gardiner, L. E., Skyner, L. J., Berkvens, C. N. and Anderson, W. G. (2014). Validation of a radioimmunoassay-based fecal corticosteroid assay for Richardson's ground squirrels *Urocitellus richardsonii* and behavioural correlates of stress. *Current Zoology*, 60(5), 591-601. <https://doi.org/10.1093/czoolo/60.5.591>
- [31]. Hogenelst, K., Soeter, M. and Kallen, V. (2019). Ambulatory measurement of cortisol: Where do we stand, and which way to follow? *Sensing and Bio-Sensing Research*, 22, 100249.
- [32]. Jepsen, S. M., Molotch, N. P., Williams, M. W., Rittger, K. E. and Sickman, J. O. (2012). Interannual variability of snowmelt in the Sierra Nevada and Rocky Mountains, United States: Examples from two alpine watersheds, *Water Resource Research*, 48, W02529, <https://doi.org/10.1029/2011WR011006>
- [33]. Jones, S. M. and Bell, K. (2004). Plasma corticosterone concentrations in males of the skink *Egernia whitii* during acute and chronic confinement and over a diel period. *Comparative Biochemistry and Physiology Part A: Molecular and Integrative Physiology*, 137(1), 105-113. <https://doi.org/10.1007/s00360-012-0656-9>
- [34]. Kuhlman, J. R., and Martin, L. B. (2010). Captivity affects immune redistribution to skin in a wild bird. *Functional Ecology*, 24(4), 830-837. <https://doi.org/10.1111/j.1365-2435.2010.01710.x>

- [35]. Larcombe, S. D., Tregaskes, C. A., Coffey, J., Stevenson, A. E., Alexander, L. G. and Arnold, K. E. (2015). Oxidative stress, activity behaviour and body mass in captive parrots. *Conservation Physiology*, 3(1). <https://doi.org/10.1093/conphys/cov045>
- [36]. Linklater, W. L., MacDonald, E. A., Flamand, J. R. B. and Czekala, N. M. (2010). Declining and low fecal corticoids are associated with distress, not acclimation to stress, during the translocation of African rhinoceros. *Animal Conservation*, 13(1), 104-111. <https://doi.org/10.1111/j.1469-1795.2009.00308.x>
- [37]. López-Olvera, J. R., Marco, I., Montané, J., Casas-Díaz, E. and Lavín, S. (2007). Effects of acepromazine on the stress response in Southern chamois (*Rupicapra pyrenaica*) captured by means of drive-nets. *Canadian Journal of Veterinary Research*, 71(1), 41.
- [38]. Marçalo, A., Pousão-Ferreira, P., Mateus, L., Duarte Correia, J. H. and Stratoudakis, Y. (2008). Sardine early survival, physical condition and stress after introduction to captivity. *Journal of Fish Biology*, 72(1), 103-120.
- [39]. Marra, P. P., Lampe, K. T. and Tedford, B. L. (1995). Plasma corticosterone levels in two species of *Zonotrichia* sparrows under captive and free-living conditions. *The Wilson Bulletin*, 296-305.
- [40]. Martin, L. B. (2009). Stress and immunity in wild vertebrates: timing is everything. *General and comparative endocrinology*, 163(1-2), 70-76. <https://doi.org/10.1016/j.ygcen.2009.03.008>
- [41]. Martin, L. B., Kidd, L., Liebl, A. L. and Coon, C. A. (2011). Captivity induces hyper-inflammation in the house sparrow (*Passer domesticus*). *Journal of Experimental Biology*, 214(15), 2579-2585. <https://doi.org/10.1242/jeb.057216>
- [42]. Mathies, T., Felix, T. A. and Lance, V. A. (2001). Effects of trapping and subsequent short-term confinement stress on plasma corticosterone in the brown treesnake (*Boiga irregularis*) on Guam. *General and Comparative Endocrinology*, 124(1), 106-114.
- [43]. Miller, R., Plessow, F., Kirschbaum, C. and Stalder, T. (2013). Classification criteria for distinguishing cortisol responders from nonresponders to psychosocial stress: evaluation of salivary cortisol pulse detection in panel designs. *Psychosomatic Medicine*, 75(9), 832-840. <https://doi.org/10.1097/PSY.0000000000000002>
- [44]. Moberg, G. P. and Mench, J. A. (Eds.) (2000). *The biology of animal stress: basic principles and implications for animal welfare*. CABI. <https://doi.org/10.1079/9780851993591.0000>
- [45]. Moore, I. T. and Jessop, T. S. (2003). Stress, reproduction, and adrenocortical modulation in amphibians and reptiles. *Hormones and Behavior*, 43(1), 39-47. [https://doi.org/10.1016/S0018-506X\(02\)00038-7](https://doi.org/10.1016/S0018-506X(02)00038-7)
- [46]. Moore, M. C., Thompson, C. W. and Marler, C. A. (1991). Reciprocal changes in corticosterone and testosterone levels following acute and chronic handling stress in the tree lizard, *Urosaurus ornatus*. *General and comparative endocrinology*, 81(2), 217-226. [https://doi.org/10.1016/0016-6480\(91\)90006-R](https://doi.org/10.1016/0016-6480(91)90006-R)
- [47]. Moreno, J., Yorrio, P., Garcia-Borboroglu, P., Potti, J. and Villar, S. (2002). Health state and reproductive output in Magellanic penguins (*Spheniscus magellanicus*). *Ethology Ecology and Evolution*, 14(1), 19-28. <https://doi.org/10.1080/08927014.2002.9522758>
- [48]. Morgan, K. N. and Tromborg, C. T. (2007). Sources of stress in captivity. *Applied animal behaviour science*, 102(3-4), 262-302. <https://doi.org/10.1016/j.applanim.2006.05.032>
- [49]. Mosconi, G., Yamamoto, K., Kikuyama, S., Carotti, M., Palermo, F. and Polzonetti-Magni, A. (2006). Neuroendocrine modulation of stress response in the anuran, *Rana esculenta*. *Amphibia-Reptilia*, 27(3), 401-408.
- [50]. Möstl, E. and Palme, R. (2002). Hormones as indicators of stress. *Domestic animal endocrinology*, 23(1-2), 67-74. [https://doi.org/10.1016/S0739-7240\(02\)00146-7](https://doi.org/10.1016/S0739-7240(02)00146-7)
- [51]. Narayan, E. J., Cockrem, J. F. and Hero, J. M. (2012). Effects of temperature on urinary corticosterone metabolite responses to short-term capture and handling stress in the cane toad (*Rhinella marina*). *General and Comparative Endocrinology*, 178(2), 301-305. <https://doi.org/10.1016/j.ygcen.2012.06.014>
- [52]. Narayan, E. and Hero, J. M. (2011). Urinary corticosterone responses and haematological stress indicators in the endangered Fijian ground frog (*Platymantis vitiana*) during transportation and captivity. *Australian Journal of Zoology*, 59(2), 79-85. <https://doi.org/10.1071/ZO11030>
- [53]. Nikoo, M., Falahatkar, B., Alekhorshid, M., Nematdost Haghi, B., Asadollahpour, A., Zarei Dangsareki, M. and Faghani, H. (2010). Physiological stress responses in kutum *Rutilus frisii* kutum subjected to captivity. *International Aquatic Research*, 2(1), 55-60.

- [54]. O'connor, T. M., O'halloran, D. J. and Shanahan, F. (2000). The stress response and the hypothalamic-pituitary-adrenal axis: from molecule to melancholia. *The Quarterly Journal of Medicine*, 93(6), 323-333. <https://doi.org/10.1093/qjmed/93.6.323>
- [55]. Olsen, D. E. and Seabloom, R. W. (1973). Adrenocortical response to captivity in *Microtus pennsylvanicus*. *Journal of Mammalogy*, 54(3), 779-781. <https://doi.org/10.2307/1378983>
- [56]. Palme, R. (1997). Measurement of cortisol metabolites in faeces of sheep as a parameter of cortisol concentration in blood. *Mammal Biology*, 62, 192-197.
- [57]. Perogamvros, I., Keevil, B. G., Ray, D. W. and Trainer, P. J. (2010). Salivary cortisone is a potential biomarker for serum free cortisol. *The Journal of Clinical Endocrinology and Metabolism*, 95(11), 4951-4958. <https://doi.org/10.1210/jc.2010-1215>
- [58]. Piersma, T. and Ramenofsky, M. (1998). Long-term decreases of corticosterone in captive migrant shorebirds that maintain seasonal mass and moult cycles. *Journal of Avian Biology*, 97-104. <https://doi.org/10.2307/3677186>
- [59]. Piersma, T., Reneerkens, J. and Ramenofsky, M. (2000). Baseline corticosterone peaks in shorebirds with maximal energy stores for migration: a general preparatory mechanism for rapid behavioral and metabolic transitions? *General and comparative endocrinology*, 120(1), 118-126. <https://doi.org/10.1006/gcen.2000.7543>
- [60]. Pifarré, M., Valdez, R., González-Rebeles, C., Vázquez, C., Romano, M. and Galindo, F. (2012). The effect of zoo visitors on the behaviour and faecal cortisol of the Mexican wolf (*Canis lupus baileyi*). *Applied Animal Behaviour Science*, 136(1), 57-62. <https://doi.org/10.1016/j.applanim.2011.11.015>
- [61]. Posthuma-Trumpie, G. A., Korf, J. and van Amerongen, A. (2009). Lateral flow (immuno) assay: its strengths, weaknesses, opportunities and threats. A literature survey. *Analytical and bioanalytical chemistry*, 393(2), 569-582. <https://doi.org/10.1007/s00216-008-2287-2>
- [62]. Quispe, R., Villavicencio, C. P., Addis, E., Wingfield, J. C. and Vasquez, R. A. (2014). Seasonal variations of basal cortisol and high stress response to captivity in *Octodon degus*, a mammalian model species. *General and Comparative Endocrinology*, 197, 65-72. <https://doi.org/10.1016/j.ygcen.2013.12.007>
- [63]. Rangel-Negrín, A., Alfaro, J. L., Valdez, R. A., Romano, M. C. and Serio-Silva, J. C. (2009). Stress in Yucatan spider monkeys: effects of environmental conditions on fecal cortisol levels in wild and captive populations. *Animal Conservation*, 12(5), 496-502. <https://doi.org/10.1111/j.1469-1795.2009.00280.x>
- [64]. Rich, E. L. and Romero, L. M. (2005). Exposure to chronic stress downregulates corticosterone responses to acute stressors. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology*, 288(6), R1628-R1636. <https://doi.org/10.1152/ajpregu.00484.2004>
- [65]. Romero, L. M. (2004). Physiological stress in ecology: lessons from biomedical research. *Trends in ecology and evolution*, 19(5), 249-255. <https://doi.org/10.1016/j.tree.2004.03.008>
- [66]. Sapolsky, R. M., Romero, L. M. and Munck, A. U. (2000). How do glucocorticoids influence stress responses? Integrating permissive, suppressive, stimulatory, and preparative actions. *Endocrine reviews*, 21(1), 55-89. <https://doi.org/10.1210/er.21.1.55>
- [67]. Sheriff, M. J., Dantzer, B., Delehanty, B., Palme, R. and Boonstra, R. (2011). Measuring stress in wildlife: techniques for quantifying glucocorticoids. *Oecologia*, 166(4), 869-887. <https://doi.org/10.1007/s00442-011-1943-y>
- [68]. Siebert, U., Pozniak, B., Hansen, K. A., Nordstrom, G., Teilmann, J., Van Elk, N. and Dietz, R. (2011). Investigations of Thyroid and Stress Hormones in Free-Ranging and Captive Harbor Porpoises (*Phocoena phocoena*): A Pilot Study. *Aquatic mammals*, 37(4). <https://doi.org/10.1578/AM.37.4.2011.443>
- [69]. Stead-Richardson, E., Bradshaw, D., Friend, T. and Fletcher, T. (2010). Monitoring reproduction in the critically endangered marsupial, Gilbert's potoroo (*Potorous gilbertii*): preliminary analysis of faecal oestradiol-17 β , cortisol and progestagens. *General and Comparative Endocrinology*, 165(1), 155-162. <https://doi.org/10.1016/j.ygcen.2009.06.009>
- [70]. Steyn, D. G. (1975). The effects of captivity stress on the blood chemical values of the chacma baboon (*Papio ursinus*). *Laboratory Animals*, 9(2), 111-120. <https://doi.org/10.1258/002367775780994637>
- [71]. Suleman, M. A., Wango, E., Sapolsky, R. M., Odongo, H. and Hau, J. (2004). Physiologic manifestations of stress from capture and restraint of free-ranging male African green monkeys (*Cercopithecus aethiops*). *Journal of Zoo and Wildlife Medicine*, 35(1), 20-24. <https://doi.org/10.1638/01-025>

- [72]. Sykes, K. L. and Klukowski, M. (2009). Effects of acute temperature change, confinement and housing on plasma corticosterone in water snakes, *Nerodia sipedon* (Colubridae: Natricinae). *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology*, 311(3), 172-181. <https://doi.org/10.1002/jez.515>
- [73]. Titon, S. C. M., Assis, V. R., Titon Junior, B., Cassettari, B. D. O., Fernandes, P. A. C. M. and Gomes, F. R. (2017). Captivity effects on immune response and steroid plasma levels of a Brazilian toad (*Rhinella schneideri*). *Journal of Experimental Zoology Part A: Ecological and Integrative Physiology*, 327(2-3), 127-138. <https://doi.org/10.1002/jez.2078>
- [74]. Titon, S. C. M., Junior, B. T., Assis, V. R., Kinker, G. S., Fernandes, P. A. C. M. and Gomes, F. R. (2018). Interplay among steroids, body condition and immunity in response to long-term captivity in toads. *Scientific reports*, 8(1), 1-13. <https://doi.org/10.1038/s41598-018-35495-0>
- [75]. Trumble, S. J., O'Neil, D., Cornick, L. A., Gulland, F. M., Castellini, M. A. and Atkinson, S. (2013). Endocrine Changes in Harbor Seal (*P hoca vitulina*) Pups Undergoing Rehabilitation. *Zoo Biology*, 32(2), 134-141. <https://doi.org/10.1002/zoo.21036>
- [76]. Tyrrell, C. and Cree, A. (1994). Plasma corticosterone concentrations in wild and captive juvenile *tuatara* (*Sphenodon punctatus*). *New Zealand journal of zoology*, 21(4), 407-416. <https://doi.org/10.1080/03014223.1994.9518010>
- [77]. Vera, F., Antenucci, C. D. and Zenuto, R. R. (2011). Cortisol and corticosterone exhibit different seasonal variation and responses to acute stress and captivity in tuco-tucos (*Ctenomys talarum*). *General and comparative endocrinology*, 170(3), 550-557. <https://doi.org/10.1016/j.ygcen.2010.11.012>
- [78]. West, D. B. and York, B. (1998). Dietary fat, genetic predisposition, and obesity: lessons from animal models. *The American Journal of clinical nutrition*, 67(3), 505S-512S. <https://doi.org/10.1093/ajcn/67.3.505S>
- [79]. Wingfield, J. C., Smith, J. P. and Farner, D. S. (1982). Endocrine responses of white-crowned sparrows to environmental stress. *The Condor*, 84(4), 399-409. <https://doi.org/10.2307/1367443>
- [80]. Yang, L., Wang, W., Huang, S., Wang, Y., Wronski, T., Deng, H. and Lu, J. (2019). Individual stress responses of white rhinoceros (*Ceratotherium simum*) to transport: implication for a differential management. *Global Ecology and Conservation*, 17, e00588. <https://doi.org/10.1016/j.gecco.2019.e00588>
- [81]. Zerani, M., Amabili, F., Mosconi, G. and Gobetti, A. (1991). Effects of captivity stress on plasma steroid levels in the green frog, *Rana esculenta*, during the annual reproductive cycle. *Comparative Biochemistry and Physiology Part A: Physiology*, 98(3-4), 491-496. [https://doi.org/10.1016/0300-9629\(91\)90436-G](https://doi.org/10.1016/0300-9629(91)90436-G)

HOW TO CITE THIS ARTICLE?

MLA

Islam, P. et al. "Stress Biomarker: A Review on Glucocorticoids Concentration Pattern & its Impact on Captive Animals". *Journal of Fisheries, Livestock and Veterinary Science* 02(02) (2022): 88-100.

APA

Islam, P., Saha, P., Khatun, P., Hossain, M. I., Islam, A. and Sachi, S. (2022). Stress Biomarker: A Review on Glucocorticoids Concentration Pattern & its Impact on Captive Animals. *Journal of Fisheries, Livestock and Veterinary Science*, 02(02), 88-100.

Chicago

Islam, P., Saha, P., Khatun, P., Hossain, M. I., Islam, A. and Sachi, S. "Stress Biomarker: A Review on Glucocorticoids Concentration Pattern & its Impact on Captive Animals". *Journal of Fisheries, Livestock and Veterinary Science* 02(02) (2022): 88-100.

Harvard

Islam, P., Saha, P., Khatun, P., Hossain, M. I., Islam, A. and Sachi, S. 2022. Stress Biomarker: A Review on Glucocorticoids Concentration Pattern & its Impact on Captive Animals. *Journal of Fisheries, Livestock and Veterinary Science*, 02(02), pp. 88-100.

Vancouver

Islam, P, Saha, P, Khatun, P, Hossain, MI, Islam, A and Sachi, S. Stress Biomarker: A Review on Glucocorticoids Concentration Pattern & its Impact on Captive Animals. *Journal of Fisheries, Livestock and Veterinary Science*. 2022 August 02(02):88-100.