

Published with Open Access at **Journal BiNET**

Vol. 22, Issue 01: 1829-1834

Journal of Bioscience and Agriculture ResearchJournal Home: www.journalbinet.com/jbar-journal.html

Response of groundnut (*Arachis hypogea*) to the exogenous application of growth hormones (IAA and GA₃)

Mukundraj B. Patil

Dept. of Botany, Late Ramesh Warpudkar Arts, Commerce and Science College, Sonpeth, Dist. Parbhani (MS), India.

✉ Corresponding Author: mukundrajpatil@gmail.com.

Article received: 13.07.19; Revised: 20.09.19; First published online: 25 October 2019.

ABSTRACT

Experiments were carried out to study the effect of Indole Acetic Acid (IAA) and Gibberellic Acid (GA₃) on groundnut (TAG-24). In this study, the effect of growth regulators was studied on morphological as well as yield parameters of groundnut plant. The sowing of seed was carried out in eleven plots of 1m x 1m, one for control and with 10ppm, 20ppm, 30ppm, 40ppm and 50ppm concentrations of IAA and GA₃. Different concentrations of IAA and GA₃ were applied twice 30 and 60 days after sowing. Compared to IAA, GA₃ hr hormone was found more effective to increase the height of the plant, the number of root nodules, the size of the seeds, number of pods, weight of seeds as well as the yield of groundnut. IAA has affected all the parameters, but except the height of the plant and length of the root, all the parameters showed non-significant change. Exogenous application of GA₃ enhanced the number of root nodules and number of pods per plant, but it was found reduced due to the application of IAA. Thus, application of GA₃ (i.e. GA₃ at 50ppm) can be suggested for higher yield of groundnut.

Key Words: IAA, GA₃, Root nodules, Morphological characters and Groundnut yield

Cite Article: Patil, M. B. (2019). Response of groundnut (*Arachis hypogea*) to the exogenous application of growth hormones (IAA and GA₃). Journal of Bioscience and Agriculture Research, 22(01), 1829-1834. **Crossref:** <https://doi.org/10.18801/jbar.220119.225>



Article distributed under terms of a Creative Common Attribution 4.0 International License.

I. Introduction

Indian economy is greatly influenced by various agricultural activities, production and agribusiness. As India has different of geographical and climatic conditions, hence a variety of crops are cultivated and grown year-round. In India, oilseeds contribute 14% of the total area under cultivation (GOI, 2015). India is the fourth largest edible oil economy in the world and contributes about 10% of the world oilseed production. Groundnut is one of the major nine oilseeds in India. Groundnut is one of our country's largest cash crops. It is a commodity that is low-priced but a precious source of all nutrients. But the area under cultivation and production of groundnut was decreased during 2015-16 periods due to shortage of rainfall and drought conditions. There were nine states with more than 100 thousand tons of groundnut production; in 2015-16, Gujarat, Rajasthan, Tamil Nadu, Andhra Pradesh, Karnataka, Madhya Pradesh, Maharashtra, Telangana and West Bengal (OGD PMU Team 2017). However, compared to the world scenario yield of groundnut is very less (GOI 2015) in India. Hence, efforts should

be made to increase the yield to attract farmers towards the cultivation of groundnut. Growth hormones can contribute to the improved growth and development of field crops with higher yields, e.g. groundnut (Khan et al. 2011). In addition, auxins and GA₃ are important growth hormones that can affect the development of plant morphological and yield contributing attributes. Besides their primary role in growth regulation, growth hormones also affect timing of flowering, the sex of the flowers, senescence of leaves and fruits, fruit development, etc. Most of the workers have studied the effect of growth regulators on seed germination (Chauhan et al. 2009; Patil et al. 2012; Dhoran and Gudadhe, 2012; Patil and Bhosle, 2017). Recent studies (Emonger and Ndambole, 2011; Rastogi et al. 2013; Khairul Mazed, et al. 2015) have focused on the impact of growth regulators on yield contributing parameters, hence it can increase the yield of the crop. This present study deals with the response of groundnut to the IAA and GA₃ in relation to morphological and yield contributing attributes, such as height of plant, number of root nodules, number of pods per plant, seed size, seed weight and yield.

II. Materials and Methods

The experiment was carried out in the Department of Botany, Late Ramesh Warpudkar Arts, Commerce and Science College, Sonpeth, Parbhani (MS) district, India, from December 2013 to March 2014 to study effect of IAA and GA₃ on morphological and yield parameters of Groundnut (TAG-24). A Stock solution of 100 ppm GA₃ and IAA was prepared by dissolving 50mg into 500 ml of water (Before preparation of Stock solution GA₃ was dissolved in small quantity of alcohol). It was diluted to prepare different concentrations of growth hormones using $N_1V_1=N_2V_2$ formula to prepare 10ppm, 20ppm, 30ppm, 40ppm and 50ppm IAA and GA₃. To study effect of growth hormones on growth and yield parameters plots of 1meter X 1meter were prepared. Groundnut (TAG-24) seeds were sowed in these plots. Growth hormones were applied twice first 30 days after and second 60 days after date of sowing. Spraying of growth hormones was carried out early in the morning with hand spray. The height of the plant and length of the root was measured physically with help of Scale in centimeter. The number of root nodules and the Number of pods per plant were counted numerically. Each treatment was replicated three times and average values were tabulated in Table 1.

Size of seed: Size of the seed was determined by pouring seeds into the 25 ml measuring cylinder as number of seeds per 25ml increases size of the seed decreases and vice-versa.

Weight of seed: After harvesting, 100 seeds were selected randomly from each plot and they were weighed using electronic balance and represented in gram.

Yield: At the end of the season, crop from each plot of 1x1m was harvested and threshed separately and weighed using the electronic balance in kilograms. It was multiplied by 10000 and converted into kg/hectare.

Statistical analysis: The mean, standard deviation (SD) and coefficient of variation (CV) have been calculated as described by Mungikar (2003). Standard error (SE) was calculated as $S.D. / \sqrt{n}$ (n = number of observations), and Critical difference (CD) was calculated as SE multiplied with t value for n^{-1} . (C.D.= S.E. x t value for n^{-1}).

III. Results and Discussion

In control, the average height of the groundnut plant was 35cm while maximum height of the plant was recorded in plants treated with 50ppm IAA which was 40.2cm (Table 01). Application of IAA and GA₃ has shown increase in the height of the plant. This increase in the height of the plant was highly significant in all treatments. The same results were found by Bora and Sarma (2006), Rahman et al. (2004) and Patil (2017) to increase plant height due to application of plant growth hormones. Different concentrations of the IAA and GA₃ have shown significant increase in the length of roots. The average length of roots in control was 20cm (Table 01). Application of GA₃ increased length of roots up to 30ppm concentration it was decreased at higher concentrations of GA₃ (40 and 50ppm). There was direct proportion between the concentration of IAA and length of root.

An increase in the root length was statistically significant at $p=0.01$ in all concentrations of IAA and in 30ppm GA₃. In 10 and 20ppm IAA it was significant at $p=0.05$. The concentration of Gibberellic acid has

shown direct effect on the number of root nodules in the groundnut. In control the number of root nodules per plant was 28 which were increased up to 38 in plants treated with 50ppm GA₃(Table 01). An increase in the number of nodules per plant in 10ppm and 20ppm GA₃ was statistically non-significant while in 30ppm 40ppm and 50ppm GA₃ increase in the number of nodules was statistically significant at p=0.001. Plants treated with different concentrations of IAA have shown variable effect on the number of root nodules per plant. 10 and 20 ppm concentration of IAA increases but this increase in the number of nodules is statistically non-significant. It was decreased (26 per plant) than control (28/plant) in the plants treated with 40 and 50ppm IAA but this decrease was statistically non-significant. Such type of increased nodulation was recorded by [Dobert \(1992\)](#), [Lievens \(2005\)](#) and [Emonger and Ndambole \(2011\)](#) due to the application of Growth Hormones. While the results obtained by [Williams and Mallorca \(1984\)](#) in Soybean were contrast with the results obtained here.

The number of pods per plant was constantly increased with the concentration of GA₃. Increased number of pods due to application 10ppm GA₃ is significant at p=0.05 while other four concentrations of GA₃ have shown increase in the number of pods which was significant at p=0.01. The maximum number of pods per plant was recorded in plants applied with 50ppm GA₃ that was 38 pods per plant. The role of IAA to increase number of pods in groundnut is very little. The number of pods per plant was increased up to 30ppm IAA concentration which was non-significant statistically. The plants applied with 40ppm IAA show no increase in the number of pods while 50ppm IAA shows reduction in the number of pods per plant (23 pods per plant) this decrease is also statistically non-significant. The same results were found by [Emonger \(2007\)](#) in Cowpea, [Resmi and Gopalkrishnan \(2004\)](#) with NAA (15, 30, and 45 ppm) and 2-4D (2ppm) in *Vigna unguiculata*. [Rathod et al. \(2015\)](#) recorded maximum number of green pods per plant in French bean due to application of 100 ppm GA₃.

The size of the seed was influenced only by higher concentration of GA₃. 10ppm GA₃, as well as all the concentrations of IAA, has no significant effect on size of the seed. 20ppm GA₃ has shown significant increase in the size of groundnut seeds it is significant at p=0.05, while 30, 40 and 50 ppm GA₃ has maximum seed size which is significant at p=0.01.

The weight of 100 seeds of groundnut in control was 34.9gm. Only 40ppm and 50ppm GA₃ was effective to increase weight of the seeds. A maximum weight of 100 seeds was found in the plants applied with 50ppm GA₃ (39.2gm). The increase in the weight of 100 seeds in the plant applied with 40 and 50ppm GA₃ was statistically significant at p=0.01 lower concentrations of GA₃ had shown increase in the seed weight which was non-significant. IAA did not show any significant change in the weight of 100 seeds compared to control.

Increase in the weight of seeds was also recorded by earlier workers like [Ghodrat et al. \(2012\)](#) in *Zea mays* due to application of IBA and GA₃ in 4-6 leaf stage, [Emonger \(2007\)](#) in Tswana Cowpea due to application of GA₃ and [Patil \(2017\)](#) in sunflower due to application of IAA and GA₃. Growth hormones had no same results in all kinds of plants [Ginnakoula et al. \(2012\)](#) in Lentil recorded decrease in 1000-seeds weight was by 26% in plots applied with GA₃.

It was found that 50ppm GA₃ was statistically significant at the p=0.01 level while 40ppm GA₃ was statistically significant at p=0.05 level to increase yield of Groundnut. Besides these two treatments, any other treatment was not significant statistically to enhance Yield of Groundnut. In control, yield was 1500 kg/hectare while it was maximum in 50ppm GA₃ applied plants that were 2250kg/hectare followed by 50ppm GA₃ where it was 1670 kg/hectare. Many workers have worked on this parameter and found that growth hormones are able to increase yield of the crop. [Resmi and Gopalkrishnan \(2004\)](#), [Emonger \(2007\)](#) and [Rastogi et al. \(2013\)](#) have recorded increased yield in Long bean, Cowpea, and Linseed respectively. [Ernst et al. \(2016\)](#) and [Patil \(2017\)](#) have also recorded increase in the sunflower yield due to application of Gibberellic acid. [Sarkar et al. \(2002\)](#) and [Agawane and Parhe \(2015\)](#) found increased yield in soybean due to 100 ppm IAA and 100 ppm GA₃ respectively.

Table 01. Influence of growth hormones morphological and yield parameters of groundnut

Treatments	Height of plant(cm)	Length of Root(cm)	No. of root nodules	Size of seed (seed/25ml)	Number of pods	100 seeds Weight (gm)	Yield (kg/ha)
Control	35	20.0	28.0	38.3	24.0	34.9	1500.0
GA 10ppm	37.1**	20.3*	30.0	37.7	28.0*	35.0	1500.0
GA 20ppm	37.2**	20.3*	30.0	37.3*	32.0**	35.5	1550.0
GA 30ppm	38.1**	20.4**	33.0**	36.3**	33.0**	35.7	1510.0
GA 40ppm	37.8**	19.9	34.0**	35.0**	34.0**	37.3**	1670.0*
GA 50ppm	37.6**	19.8	38.0**	35.0**	38.0**	39.2**	2250.0**
IAA 10ppm	35.8**	20.4**	30.0	38.3	26.0	34.8	1480.0
IAA20ppm	36.2**	20.6**	30.0	38.7	25.0	34.8	1510.0
IAA30ppm	37.4**	21.0**	28.0	38.3	25.0	35.0	1610.0
IAA40ppm	38.9**	21.0**	26.0	38.0	24.0	34.9	1520.0
IAA50ppm	40.2**	21.1**	26.0	38.3	23.0	35.0	1490.0
<i>Mean</i>	<i>37.4</i>	<i>20.4</i>	<i>30.3</i>	<i>37.4</i>	<i>28.4</i>	<i>35.7</i>	<i>1599.1</i>
<i>S.D.</i>	<i>1.4</i>	<i>0.4</i>	<i>3.4</i>	<i>1.3</i>	<i>4.8</i>	<i>1.3</i>	<i>213.0</i>
<i>C.V.</i>	<i>5.5</i>	<i>2.1</i>	<i>11.3</i>	<i>3.4</i>	<i>17.0</i>	<i>3.7</i>	<i>13.3</i>
<i>S.E.</i>	<i>0.4</i>	<i>0.1</i>	<i>1.0</i>	<i>0.4</i>	<i>1.5</i>	<i>0.4</i>	<i>64.2</i>
<i>C.D. 5 %</i>	<i>1</i>	<i>0.3</i>	<i>2.3</i>	<i>0.9</i>	<i>3.2</i>	<i>0.9</i>	<i>143.2</i>
<i>C.D. 1 %</i>	<i>1.4</i>	<i>0.4</i>	<i>3.3</i>	<i>1.2</i>	<i>4.6</i>	<i>1.3</i>	<i>203.6</i>

(* Significant at p=0.05; ** Significant at p=0.01).

IV. Conclusion

From the above experiment, it is evident that IAA and GA₃ both are playing very important role to increase height of the groundnut plant while GA₃ can be applied to increase number of root nodules, number of pods per plant, seed size, seed weight and consequently the higher yield of groundnut.

Acknowledgments

The author is thankful to UGC (WRO Pune) for financial assistance under Minor Research Project.

Reference

- (1) Agawane, R. B. and Parhe, S. D. (2015). Effect of seed priming on crop growth and seed yield of soybean (*Glycine max* (L) Merrill). *The Bioscan*, 10, 265-270.
- (2) Bora, R. K. and Sarma, C. M. (2006). Effect of gibberellic acid and cycocel on growth, yield and protein content of pea. *Asian Journal of Plant Sciences*. 5(2), 324-330.
<https://doi.org/10.3923/ajps.2006.324.330>
- (3) Chauhan, J. S. Tomar, Y. K. Badoni Anoop, N. Indrakumar Singh, Seema Ali, Debarati, Rawat A. S. and Nautiyal V. P. (2009). Morphology and Influence of Various Plant Growth Substances on Germination and Early Seedling Growth in *Macrotyloma uniflorum* (Lam.). *Journal of American Science*, 5(6), 43-50.
- (4) Dhoran, V. S. and Gudadhe, S. P. (2012). Effect of plant growth regulators on seed germination and seedling vigor in *asparagus sprengeri regelin*. *International Research Journal of Biological Sciences*, 1(7), 6-10.
- (5) Dobert, R. C. Rood, S. B. & Blevins, D. G. (1992). Gibberellins and legume-Rhizobium symbiosis: endogenous gibberellins of Lima bean (*Phaseolus lunatus* L.) stems and nodules. *Plant Physiology*, 98, 221-224. <https://doi.org/10.1104/pp.98.1.221>
- (6) Emonger, V. (2007). Gibberellic Acid (GA₃) Influence on vegetative growth, nodulation and yield of cowpea (*Vigna unguiculata* L.) Walp. *Journal of Agronomy*, 6(4), 509-517.
<https://doi.org/10.3923/ja.2007.509.517>

- (7) Emongor V. E. and Ndambole C. M. (2011). Effect of gibberellic acid on performance of cowpea. African Crop Science Conference Proceedings, Vol. 10. pp. 87 - 92.
- (8) Ernst, D. Marek, K. and Ivan, C. (2016). Effect of two different plant growth regulators on production traits of sunflower. Journal of Central European Agriculture, 17(4),998-1012.
<https://doi.org/10.5513/JCEA01/17.4.1804>
- (9) Ghodrat, V. Mohammad, J. R. Mohammad, S. T. and Abas, K. (2012). Yield and yield components of corn (*Zea mays* L.) In: Response to foliar application with indole butyric acid and gibberellic acid. American-Eurasian J. Agric. and Environ. Sci. 12(9), 1246-1251.
- (10) Giannakoula, A. E., Ilias, F. I., Maksimovic, J. J. D., Maksimovic, V. M., Zivanovic, B. D. (2012). The effects of plant growth regulators on growth, yield, and phenolic profile of lentil plants. Journal of Food Composition and Analysis. 28, 46–53.
<https://doi.org/10.1016/j.jfca.2012.06.005>
- (11) GOI- Ministry of Agriculture (2015). Agricultural statistics at glance 2014. New Delhi: Oxford University press.
- (12) Khairul Mazed, H. E. M. Akand, M. H. Israt, J. I. Jannatul, F. M. and Rahman, M. H. (2015). Effect of gibberellic acid on the growth and yield of cabbage (*Brassica oleraceavar capitata* L.). International Journal of Applied Research. 1(4), 24-29.
- (13) Khan, A. Bakht, J. E. H. A. N. Bano, A. and Malik, N. J. (2011). Effect of plant growth regulators and drought stress on groundnut (*Arachis hypogaea* L) genotypes. Pak J Bot, 43, 2397-2402.
- (14) Lievens, S. Goormachtig, S. Herder, J. D. Capoen, W. Mathis, R. Hedden, P. and Holsters, M. (2005). Gibberellins are involved in nodulation of *Sesbania rostrata*. Plant Physiology. 139, 1366-1379. <https://doi.org/10.1104/pp.105.066944>
- (15) Patil, M. B. (2017). Influence of growth hormones (IAA and GA₃) on morphological and yield parameters of sunflower (*Helianthus anus* L.). Multilogic in Science, 7(25), 175-176(b).
- (16) Mungikar, A. M. (2003). Biostatistical Analysis. Saraswati Publ. Printing press, Aurangabad (M.S.) India.
- (17) OGD PMU Team (2017). Groundnut production of various states in 2015-16. Groundnut production of various states in 2015-16. Retrieved 15 September 2018 from <https://community.data.gov.in/groundnut-production-of-various-states-in-2015-16/>.
- (18) Patil, J. G. Ahire, M. L. and Nikam, T. D. (2012). Influence of plant growth regulators on in vitro seed germination and seedling development of *Digitalis purpurea* L. The Asian and Australasian Journal of Plant Science and Biotechnology, 6 (Special Issue 1), 12-18.
- (19) Patil, M. B. and Shailaja, B. B. (2017). Effect of GA₃ on germination of certain oilseeds. Bioscience Discovery, 8(3), 483-485.
- (20) Rahman, M. S. Islam, M. N. Tahar, A. and Karim, M. A. (2004). Influence of GA₃ and MH and their time of spray-on Morphology, yield contributing characters and yield of Soyabean. Asian J. Plant Science, 3, 602-609. <https://doi.org/10.3923/ajps.2004.602.609>
- (21) Rastogi, A. Ameena, S. Brij, K. M. Mrinalini, S. Rawli, P. Pratibha, M. Munna, S. and Sudhir, S. (2013). Effect of auxin and gibberellic acid on growth and yield components of linseed (*Linum usita tissimum* L.). Crop Breeding and Applied Biotechnology, 13, 136-143.
<https://doi.org/10.1590/S1984-70332013000200006>
- (22) Rathod, R. R. Gore R. V. and Bothikar P. A. (2015). Effect of growth regulators on growth and yield of french bean (*Phaseolus vulgaris* L.) Var. Arka Komal. OSR-JAVS, 8(5), 36-39.
- (23) Resmi, R. and Gopalakrishnan, T. R. (2004). Effect of plant growth regulators on the performance of yard long bean (*Vigna unguiculata* var. sesquipedalis (L.) Verdcourt). Journal of Tropical Agriculture, 42 (1-2), 55-57.
- (24) Sarkar, P. K. Haque, M. S. and Karim, M. A. (2002). Growth analysis of soybean as influenced by GA₃ and IAA and their frequency of application. Pakistan Journal of Agronomy, 1(4), 123-126.
<https://doi.org/10.3923/ja.2002.123.126>
- (25) Williams, P. M. and de Mallorca, M. S. (1984). Effect of gibberellins and the growth retardant CCC on the nodulation of soya. Plant and Soil, 77, 53-60.
<https://doi.org/10.1007/BF02182811>

HOW TO CITE THIS ARTICLE?**Crossref:** <https://doi.org/10.18801/jbar.220119.225>**MLA**

Patil, M. B. "Response of groundnut (*Arachis hypogea*) to the exogenous application of growth hormones (IAA and GA3)." *Journal of Bioscience and Agriculture Research* 22(01) (2019): 1829-1834.

APA

Patil, M. B. (2019). Response of groundnut (*Arachis hypogea*) to the exogenous application of growth hormones (IAA and GA3). *Journal of Bioscience and Agriculture Research*, 22(01), 1829-1834.

Chicago

Patil, M. B. "Response of groundnut (*Arachis hypogea*) to the exogenous application of growth hormones (IAA and GA3)." *Journal of Bioscience and Agriculture Research* 22(01) (2019): 1829-1834.

Harvard

Patil, M. B. 2019. Response of groundnut (*Arachis hypogea*) to the exogenous application of growth hormones (IAA and GA3). *Journal of Bioscience and Agriculture Research*, 22(01), pp. 1829-1834.

Vancouver

Patil, MB. Response of groundnut (*Arachis hypogea*) to the exogenous application of growth hormones (IAA and GA3). *Journal of Bioscience and Agriculture Research*. 2019 October 22(01): 1829-1834.